

---

**Range extension for the sea cucumber *Selenkothuria lubrica*  
(Holothurioidea) the west coast of Baja California Sur, Mexico**

Received 07-III-2002.

Accepted 21-III-2002.

*Selenkothuria lubrica* (Selenka, 1867). The present work records *S. lubrica* (= *Holothuria lubrica*) along the west coast of Baja California Sur to Puerto Cortés: (24°25'15.2" N and 111°51'19.9" W), San Juanico: (26°14'35.9" N and 112°28'27.9" W) at Bahía Tortugas, (27°37'09.3" N and 114°50'55.8" W). The local fishermen know them as "miones". In San Juanico, organisms are observed with body lengths from 150 to 170 mm and densities that range from 15 to 18 ind/m<sup>2</sup>. They occur in the bay La Paz, with 13 to 35 ind/m<sup>2</sup> and body lengths from 12 to 170 mm. Acknowledgments: I am grateful to Dr. Philip Lambert, M.Sc., Invertebrate Curator, from Royal British Columbia Museum, for revisions to manuscript; to M. en

C.M. Dinorah Herrero Perezrul for taxonomic species identification.

#### REFERENCES

- Brusca, R.C. 1980. Common intertidal invertebrates of the Gulf of California. The University of Arizona, Tucson, Arizona. 513 pp.
- Clark, H.L. 1923. Supplementary report on Echinoderms from Lower California. Bull. Am. Mus. Nat. Hist. 48:147-163.
- Deichmann, E. 1958. The Holothuroidea collected by the Velero III and IV during the years 1932 to 1954. Part II, Aspidochirota. Allan Hancock Pac. Exped. II(2): 314-318.
- Hendrickson, J.R. & R.C. Brusca. 1975. Biological studies in the Gulf of California. X [a collection of student research papers]. Univ. Arizona, Tucson.

---

José R. Turrubiates M.

Instituto Nacional de la Pesca, Centro Regional de Investigación Pesquera La Paz, Carretera a Pichilingue km. 1, La Paz, Baja California Sur, México; jrturrubiates@yahoo.com