

Effects of salinity on growth and survival in five *Artemia franciscana* (Anostraca: Artemiidae) populations from Mexico Pacific Coast

Jorge Castro-Mejía¹, Talía Castro-Barrera¹, Luis Héctor Hernández-Hernández², José Luis Arredondo-Figueroa³, Germán Castro-Mejía¹ & Ramón de Lara-Andrade¹

1. Laboratorio Alimento Vivo, Departamento El Hombre y su Ambiente, Universidad Autónoma Metropolitana-Xochimilco, Calzada del Hueso No.1100, Colonia Villa Quietud, Distrito Federal, 04960, México, Tel 5255-5483 7151; camj7509@correo.xoc.uam.mx, cabt7515@correo.xoc.uam.mx, gecastromejia@correo.xoc.uam.mx, rlara@correo.xoc.uam.mx
2. Laboratorio de Producción Acuícola (Acuario), UNAM FES Iztacala, Av. de los Barrios 1, Los Reyes Iztacala Tlalnepantla, Estado de México, 54090, México, Tel 5255-56231197; luish3@yahoo.com
3. Planta Experimental de Producción Acuícola, Universidad Autónoma Metropolitana-Iztapalapa, Avenida San Rafael Atlixco No. 186, Colonia Vicentina, Distrito Federal, 09340, México, Tel 5255-58046585; afjl@xanum.uam.mx

Received 23-XI-2009.

Corrected 02-VIII-2010.

Accepted 01-IX-2010.

Abstract: Salinity is an important factor influencing growth and survival of aquatic organisms such as *Artemia*, a valuable aquaculture species. This study evaluated the effects of salinity on *A. franciscana* populations from different water bodies in Mexico's Pacific Coast. With this purpose, five autochthonous bisexual *Artemia* populations were tested to assess their survival and growth values against salinities of 40, 60, 80, 100 and 120g/l, under laboratory conditions (25±2°C; pH 8-10; constant light and aeration). The organisms were fed with 100mL of rice bran and 2L of *Tetraselmis suecica* (500 000cel/ml). The culture experiments were made in 200L plastic tanks, and survival and growth final values were obtained after 21 culture days. Survival and growth curves were determined by a regression analysis (R²). The significant differences between salinities were determined by ANOVA test (p<0.05). The best survival and growth rates were found at salinities of 100-120g/l. When the Mexican *Artemia* populations were cultivated at 40g/l of salinity, 100% mortality was observed in the juvenile stage. This study determined that survival and growth values of *A. franciscana* populations increased with salinity. The five *A. franciscana* populations presented significant differences in their survival rate under various salinity regimes. The studied populations experienced high mortality at salinities under 60g/l and over 200g/l, and especially during the metanauplius stage. The present study confirms that growth rates in Mexican *A. franciscana* populations from Pacific Coast habitats are not inversely proportional to salinity. These *A. franciscana* populations should be cultured at 100-120g/l of salinity to obtain better survival and growth rates. This data is useful to improve culture systems in aquaculture biomass production systems. Rev. Biol. Trop. 59 (1): 199-206. Epub 2011 March 01.

Key words: *Artemia franciscana*, salinity, growth, survival, Mexico, Pacific Coast.

The genus *Artemia* (Crustacea: Anostraca) consists of bisexual and parthenogenetic species that are morphologically and biometrically similar. A group named "New World" species is composed of *A. franciscana* (Kellogg 1906) (North, Central and South America), *A. persimilis* (Piccinelli & Prosdocimi 1968) (Argentina) and *A. monica* (Verrill 1869) (USA); another

group named "Old World" species is represented by *A. salina* (Leach 1819) (Mediterranean basin), *A. urmiana* (Günther 1900) (Urmia lake, Iran), *A. sinica* (Cai 1989) (China and neighboring countries), *A. tibetiana* (Abatzopoulos *et al.* 1998) (Tibet's high Plateau), *Artemia* sp. (Pilla & Beardmore 1994) (Kazakhstan) and parthenogenetic forms located only in the Old

World, i.e., *A. parthenogenetica* (Abatzopoulos *et al.* 2002). The genus *Artemia* comprises a complex of sibling species and superspecies defined by a criterion of reproductive isolation (Browne & Bowen 1991).

These organisms are largely distributed in inland and coastal hypersaline body waters (Triantaphyllidis *et al.* 1998, Castro *et al.* 2000, Van Stappen 2002, El-Bermawi *et al.* 2004). *Artemia* may inhabit chloride, sulphate or carbonate waters and combinations of more than two anions (Bowen *et al.* 1985, Lenz 1987). *Artemia* are among the few organisms that can adapt to very diverse living conditions, including salinities as low as 10g/L (Abatzopoulos *et al.* 2006a, Abatzopoulos *et al.* 2006b) and high as 340g/L (Post & Youssef 1977).

Some information is available on the survival and growth rate characteristics of bisexual and parthenogenetic *Artemia* populations (Vanhaecke *et al.* 1984, Browne *et al.* 1984, Wear & Haslett 1986, Browne & Bowen 1991, Triantaphyllidis *et al.* 1995, Triantaphyllidis *et al.* 1997a, Triantaphyllidis *et al.* 1997b, Browne & Wanigasekera 2000, Abatzopoulos *et al.* 2003, Baxevanis *et al.* 2004, Castro 2004, El-Bermawi *et al.* 2004, Abatzopoulos *et al.* 2006b, Agh *et al.* 2008) cultivated in different

salinities. Most of these studies mentioned morphometry, life span and reproductive characteristics and also contributed to the evaluation of genetic and environmental components of variance in sexual and clonal *Artemia*. However, the importance of salinity as a principal factor modifying growth and survival of world *Artemia* populations and the effects of the salinity on *A. franciscana* populations from Mexico's Pacific Coast have been little studied, if at all.

The aim of this study was to determine the effects of salinity on the survival and growth values of five Pacific Coast Mexican *Artemia* populations cultured in laboratory at five salinities, to provide additional data for a better understanding of their adaptation patterns. This data can be useful for a better culture system in aquaculture biomass production.

MATERIAL AND METHODS

Populations used in the experiment:

This study was conducted at the Laboratorio de Alimento Vivo of the Universidad Autónoma Metropolitana-Xochimilco, Mexico. The locations from where the cysts were collected are listed in Table 1.

TABLE 1

List of Mexican *A. franciscana* populations studied in this work, abbreviations used and geographical locations

Site	State	Abbreviation	Geographical location
Yavaros	Sonora	YAV	26°41' N; 109°31' W
Bahía de Lobos	Sonora	LOBOS	27°18' N; 110°30' W
Ohuira	Sinaloa	OHUI	24°18' N; 107°22' W
El Marqués	Oaxaca	MARQ	16°09' N; 95°14' W
Juchitán	Oaxaca	JUCH	16°26' N; 95°01' W

Culture experiments: The cysts used in this study were stored at -10°C to keep dehydration. An amount of 0.5g of cyst from each strain were hatched at 40g/l of salinity, a pH of 8-10, and a water temperature of 24±2°C, and under conditions of constant illumination and aeration (Castro *et al.* 2003, Castro 2004). The

newly hatched nauplii were placed into separate tanks and then transferred to 200l plastic tanks with 160l of differing salinities (40, 60, 80, 100 and 120g/l). The density of the *Artemia* nauplii was adjusted to one organism per 100ml to avoid crowding and growth problems. The animals were fed 100ml of rice bran (75g/l

per 90g/l saline water) and 2l of *Tetraselmis suecica* (500 000 cells/ml). The organisms were cultured until adulthood, when it was possible to observe mating. Culture conditions maintained in plastic tanks were: temperature of 24±2°C, pH of 8-10 and dissolved oxygen of 2mgO₂/ml.

Survival rate: The survival value was determined after 21 culture days; the mean density was taken by counting ten 500ml samples. The results (expressed as percentages) were log-transformed and plotted against time (expressed in days).

Growth rate: One hundred sexually mature organisms (50 females and 50 males) were separated from each population to obtain total length (TL) biometry data. Adult organisms were preserved with drops of acetic acid and the TL was measured with a dissection microscope equipped with a camera and Image-ProPlus 7.0 (MediaCybernetics®) software program.

For data analysis, stem and leaf displays and Box Plot were performed to ensure that the assumption of normality was met for each

data set. A descriptive statistical analysis was made to obtain mean values and standard deviation for length biometry. Analyses of variance (ANOVA) were also used to determine significant differences among the five populations (Tatsuoka 1970, Kachigan 1991). The least significant differences (LSD) pairwise comparison (Tukey; p<0.05) was used to compare pairs of sample means after ANOVAs for TL character studied were generated. Type classifications were based on populations, and were grouped for biometric analysis according to the specific salinity of their culture medium (Sokal & Rohlf 1981, Kachigan 1991). The SYSTAT 10.2 (Systat Software Inc., California, USA) software package was used for the statistical analysis.

Regression analysis (Sokal & Rohlf 1981) for the survival and growth rates of each population and salinity was performed with the Excel 2003 statistical program (Microsoft Corp. Washington, USA).

RESULTS

Survival: The respective means and standard deviation values are shown in Table 2. Organisms from all populations at 40g/l

TABLE 2
Mean values ± standard deviation of survival (%) and total length (mm) of the five Mexican *A. franciscana* Pacific Coast populations studied

Populations	Salinities (g/L) used in experiment							
	60		80		100		120	
	Survival	TL	Survival	TL	Survival	TL	Survival	TL
YAV	37.62 ^a	7.178 ^a	65.22 ^a	6.542 ^a	79.31 ^a	7.907 ^a	78.76 ^a	8.019 ^a
	±4.33	±0.431	±4.22	±0.290	±6.10	±0.150	±3.40	±0.246
LOBOS	46.90 ^b	7.221 ^a	62.65 ^b	6.581 ^a	75.57 ^b	7.954 ^a	80.54 ^{a,b}	8.067 ^a
	±4.31	±0.555	±5.56	±0.224	±3.35	±0.183	±6.12	±0.398
OHUI	34.04 ^c	6.996 ^b	64.54 ^a	6.376 ^b	79.49 ^a	7.706 ^b	84.55 ^{b,c}	7.815 ^b
	±6.88	±0.468	±4.52	±0.188	±5.68	±0.133	±3.65	±0.271
MARQ	47.86 ^b	7.720 ^c	64.76 ^a	7.035 ^c	77.81 ^{a,b}	8.504 ^c	78.99 ^a	8.624 ^c
	±4.07	±0.540	±6.09	±0.178	±6.52	±0.168	±4.27	±0.366
JUCH	35.70 ^{a,c}	8.297 ^d	67.05 ^c	7.561 ^d	79.56 ^a	9.139 ^d	80.40 ^a	9.269 ^d
	±5.86	±0.439	±4.84	±0.139	±3.94	±0.155	±3.88	±0.263

Populations sharing similar letter(s) in each column are not significantly different (Tukey, p<0.05). TL=total length.

salinity died after seven culture days (during the transition period of metanauplii to the juvenile stages). All populations showed their lowest survival rates after 21 days with a salinity of 60g/l. Almost all populations showed the highest survival level at 120g/l of salinity (79-80%) (Table 2).

When survival rate at the different salinities were compared, the regression analysis showed a strong correlation ($R^2=0.98-0.99$), with a polynomial second grade regression curve (Fig. 1). The actual tendency indicates that as salinity increased, the survival rate decreased.




Fig. 1. Survival values and binomial trend line of five Pacific Coast Mexican *Artemia* populations reared in five different salinities (40, 60, 80, 100 and 120g/l).

Growth rate: The total length means \pm standard deviation values of Pacific Coast Mexican *Artemia* populations are shown in Table 2.

The populations cultivated at 80g/l salinity exhibited the lowest mean growth rate values (6.5-7.5 mm); the populations cultivated at 120g/l salinity showed the highest (7.8-9.2mm). Only the LOBO and OHUI populations did not

show significant differences (Tukey, $p < 0.05$) between all cultured salinities.

When comparing the results of growth rate expressed as total length over the experimental period, the data showed a strong correlation ($R^2=0.8513$) with a second grade polynomial curve regression and began to decrease at higher salinities or the organisms died in the juvenile stage (Fig. 2).




Fig. 2. Growth (with binomial trend line) expressed as total length of five Pacific Coast Mexican *A. franciscana* populations reared in five different salinities (40, 60, 80, 100 and 120g/l).

DISCUSSION

The survival rates of the five bisexual *A. franciscana* populations were affected by salinity. The studied populations showed higher mortalities at salinities under 60g/l and over 120g/l. It is important to mention that higher mortality rates were observed during meta-nauplii stage. At 60g/l of salinity, 34-48% of the hatched nauplii reached the adult stage. Although Mayer (2002), El-Bermawi *et al.* (2004) and Abatzopoulos *et al.* (2006a,b) pointed out that bisexual populations do not developed well on salinities between 100-120g/l, the studied Mexican populations showed a different behavior and are like populations of *A. parthenogenetica* (Litvinenko *et al.* 2007), *A. salina* (Naceur *et al.* 2009) and *A. urmiana* (Agh *et al.* 2008).

Agh *et al.* (2008) mentioned that mortality of 100% was observed in bisexual populations reared at salinities from 150 to 200g/l and the studied Mexican populations showed a similar pattern, as survival rates decreased over the salinity of 120g/l. On the other hand, Abatzopoulos *et al.* (2006b) reported lower survival values for *A. urmiana* at salinities of 35 and 50g/l, while higher survival rates were at 100, 140 and 180g/l of salinity.

The survival at high salinities, according Triantaphyllidis *et al.* (1995) and Van Stappen (2002) might be related to the *Artemia* osmoregulatory system function, which improves at salinities ranging from 100 to 120g/l. As well, Tackaert & Sorgeloos (1991) have mentioned a genetically imprinted ecological response, which allow the optimal development of brine shrimp at salinities between 100-180g/l as a way to avoid the presence of predators (Rodríguez-Almaraz *et al.* 2006).

Triantaphyllidis *et al.* (1995), El-Bermawi *et al.* (2004) and Agh *et al.* (2008) reported that growth rate is inversely proportional to salinity. Triantaphyllidis *et al.* (1995) reported significant differences in *A. parthenogenetica* growth in Tanguu (China) populations cultured at different salinities. These authors reported that a maximum growth of *A. franciscana*

was observed at 35g/l (10.16 ± 0.85 mm). In this research, these species experienced 100% mortality at 40g/l salinity, and showed better growth at 120g/l (9.269 ± 0.263 mm).

The present study confirms that growth and survival rate in these Mexican Pacific coast *Artemia* populations are inversely proportional to salinity, supporting the findings of Gilchrist (1960), Triantaphyllidis *et al.* (1995), El-Bermawi *et al.* (2004) and Agh *et al.* (2008). In our experiment, the best results are shown in salinities of 100 to 120g/l.

The present study supports findings by many studies on the negative impact of salinity values above 120 g/l in laboratory cultures (Brown & Wanigasekera 2000, Baxevanis & Abatzopoulos 2004, Baxevanis *et al.* 2004, Agh *et al.* 2008). In accordance with previous laboratory research on several *Artemia* species it was found that the optimal range for growth and survival in sexual and asexual strains, lies between 100-120g/l (Triantaphyllidis *et al.* 1995, Baxevanis *et al.* 2004, Agh *et al.* 2008). It is probably that these salinity tests values can be observed having a similar impact in *Artemia* life span characteristics and reproductive behavior (Agh *et al.* 2008), suggesting a subject for future experiments.

Thus, it could be concluded that high salinity is a limiting factor to control density, growth and survival of *Artemia* strains from different regions cultured under laboratory conditions. In other hand, this study would help in the management of the populations of *Artemia* from the Pacific zone of Mexico. The result information obtained in this study will be useful to produce the *Artemia* cultures which are widely use as live food in the nearby shrimp and fish farms located on the Mexico Pacific coastal zone.

ACKNOWLEDGMENTS

This work is part of the doctoral thesis of the first author, who is currently enrolled in the Biological Sciences Ph.D. program at the Universidad Autónoma Metropolitana.

RESUMEN

La salinidad es el factor más importante que modifica el crecimiento y la sobrevivencia de las poblaciones mexicanas de *Artemia* en el mundo. El efecto de este factor en las poblaciones costeras del Pacífico Mexicano ha sido poco estudiado. Cinco poblaciones bisexuales de *Artemia* provenientes de cuerpos de aguas costeras del Pacífico Mexicano, se estudiaron para determinar el efecto que tiene la salinidad de 40, 60, 80, 100 y 120g/l sobre la sobrevivencia y el crecimiento, en condiciones de laboratorio (25±2°C, pH 8-10; luz y aireación continua). Los experimentos de cultivo fueron realizados en tanques de plástico de 200l. Los valores finales de sobrevivencia y crecimiento fueron determinados después de 21 días de cultivo. Los organismos fueron alimentados con 100ml de salvado de arroz y 2l de *Tetraselmis suecica* (500cel/ml). Las curvas de sobrevivencia y crecimiento fueron determinadas usando un análisis de regresión (R²). Las diferencias significativas entre los valores promedio a las distintas salinidades fueron determinadas por una prueba de ANDEVA (p<0.05). Las mejores tasa de sobrevivencia y crecimiento se encontraron a la salinidad de 100-120g/l. Cuando las poblaciones mexicanas fueron cultivadas a 40 g/l de salinidad, hubo una mortalidad del 100% de los organismos en estadio juvenil. A la salinidad de 80g/l, los valores de sobrevivencia y crecimiento, fueron menores que cuando se cultivaron entre 100 y 120g/l. Las cinco poblaciones bisexuales de *A. franciscana* presentaron diferencias significativas en sus tasas de sobrevivencia bajo las salinidades estudiadas. Las tasas de sobrevivencia fueron afectadas por la salinidad. Las poblaciones estudiadas experimentaron alta mortalidad a salinidades por debajo de 60 g/l y por arriba de 200g/l. Es importante mencionar que las altas mortalidades se observaron en el estadio metanauplios. El presente trabajo confirma que las tasas de crecimiento de las poblaciones que costeras del Pacífico Mexicano de *A. franciscana* no son inversamente proporcionales a la salinidad. Las poblaciones de *A. franciscana* deben de ser cultivadas a una salinidad de 100-120g/l para obtener mejores tasas de sobrevivencia y crecimiento. Estos datos permiten realizar mejoras en los sistemas de cultivo de este crustáceo para la obtención de biomasa para la acuicultura.

Palabras clave: *Artemia franciscana*, salinidad, crecimiento, sobrevivencia, México, costas del Pacífico.

REFERENCES

Abatzopoulos, T.J., B. Zhang & P. Sorgeloos. 1998. *Artemia tibetiana*: preliminary characterization of a new *Artemia* species found in Tibet (People's Republic of China). Int. J. Salt Lake Res. 7: 41-44.

Abatzopoulos, T.J., J.A. Beardmore, J.S. Clegg & P. Sorgeloos. 2002. *Artemia*: Basic and Applied Biology. Kluwer Academic, Dordrecht, The Netherlands.

Abatzopoulos, T.J., N. El-Bermawi, C. Vasdekis, A.D. Baxevanis & P. Sorgeloos. 2003. Effects of salinity and temperature on reproductive and life span characteristics of clonal *Artemia* (International study on *Artemia*, LXVI). Hydrobiologia 492: 191-199.

Abatzopoulos, T.J., N. Agh, G. Van Stappen, S.M. Razavi-Rouhani & P. Sorgeloos. 2006a. *Artemia* sites in Iran. J. Mar. Biol. Assoc. United Kingdom. 86: 229-307.

Abatzopoulos, T.J., A.D. Baxevanis, G.V. Triantaphyllidis, G. Criel, E.L. Pador, G. Van Stappen & P. Sorgeloos. 2006b. Quality evaluation of *Artemia urmiana* Günther (Urmia Lake, Iran) with special emphasis on its particular cyst characteristics (International Study on *Artemia*, LXIX). Aquaculture 254: 442-454.

Agh, N., G. Van Stappen, G. Bossier, P. Sepehri, H. Lofti, V. Rouhani, S.M.R.P. Sorgeloos. 2008. Effects of salinity on survival, growth, reproductive and life span characteristics of *Artemia* populations from Urmia Lake and neighboring lagoons. Pak. J. Biol. Sci. 11: 164-172.

Baxevanis, A.D., N. El-Bermawi, T.J. Abatzopoulos & P. Sorgeloos. 2004. Salinity effects on maturation, reproductive and life span characteristics of four Egyptian *Artemia* populations. Hydrobiologia 513: 87-100.

Baxevanis, A.D. & T.J. Abatzopoulos. 2004. The phenotypic response of ME2 (M Embolon, Greece) *Artemia* clone to salinity and temperature. J. Biol. Res. 1: 107-114.

Bowen, S.T., E.A. Fogarino, K.N. Hitchner, G.L. Dana, V.H.S. Chow, M.R. Buoncristiani & J.R. Carl. 1985. Ecological isolation in *Artemia*: population differences in tolerance of anion concentrations. J. Crust. Biol. 5: 106-129.

Browne, R.A., S.E. Salee, D.S. Grosch, W.O. Segreti & S.M. Purser. 1984. Partitioning genetics and environmental components of reproduction and life span in *Artemia*. Ecology 65: 949-969.

Browne, R.A. & S.T. Bowen. 1991. Taxonomy and population genetics of *Artemia*, p. 221-235. In R.A. Browne, P. Sorgeloos & C.N.A. Trotman CNA (eds.). *Artemia* Biology. CRC Boca Raton, Florida, USA.

Browne, R.A. & G. Wanigasekera. 2000. Combined effects of salinity and temperature on survival and reproduction of five species of *Artemia*. J. Exp. Mar. Biol. Ecol. 244: 29-44.

Castro, M.J. 2004. Caracterización Morfométrica y reproductiva (hibridación) de siete poblaciones de *Artemia franciscana* en México, en relación a su hábitat y

- latitud. Master Thesis, Universidad Autónoma Metropolitana, México.
- Castro, B.T., A.S. Malpica, J.M. Castro, G.M. Castro & R.A. De Lara. 2000. Environmental and biological characteristics of *Artemia* ecosystems in México: and updated review, p. 191-202. *In* M. Munawar, S.G. Lawrence, I.F. Munawar & D.F. Malley (eds.). Aquatic Ecosystems of México. Status and scope. Backhuys, Leiden, The Netherlands.
- Castro, B.T., J.M. Castro & G.M. Castro. 2003. *Artemia*, p. 67-81. *In* T. Castro B. (Comp.) Alimento vivo para organismos acuáticos. AGT, México.
- El-Bermawi, N., A.D. Baxevanis, T.J. Abatzopoulos, G. Van Stappen & P. Sorgeloos. 2004. Salinity effects on survival, growth and morphometry of four Egyptian *Artemia* populations (International study on *Artemia*, LXVII). *Hydrobiologia* 523: 175-188.
- Kachigan, S.K. 1991. Multivariate statistical analysis. A conceptual introduction. Radius, New York, USA.
- Lenz, P.H. 1987. Ecological studies on *Artemia*: a review, p. 5-18. *In* P. Sorgeloos, D.A. Bengtson, W. Declair & E. Jaspers (eds.). *Artemia* Research and its Applications, Vol. 3. Ecology, Culturing, Use in Aquaculture. Universa, Wetteren, Belgium.
- Litvinenko, L.I., A.V. Koslov, A.I. Kovalenko & D.S. Bauer. 2007. Salinity of water as a factor to determine the development of the brine shrimp *Artemia* populations in Siberian lakes. *Hydrobiologia* 576: 95-101.
- Mayer, R.J. 2002. Morphology and biometry of three populations of *Artemia* (Branchiopoda: Anostraca) from the Dominican Republic and Puerto Rico. *Hydrobiologia* 486: 29-38.
- Naceur, H.B., A.B. Rejeb-Jenhani & M.S. Romdhane. 2009. Ecobiological survey of the brine shrimp *Artemia salina* from Sabkhet El Adhibet (south-east Tunisia). *J. Mar. Biol. Assoc. United Kingdom* 89: 1109-1116.
- Pilla, E.J.S. & J.A. Beardmore. 1994. Genetic and morphological differentiation in old world bisexual species of *Artemia* (the brine shrimp). *Heredity* 72: 47-56.
- Post, F.J. & N.N. Youssef. 1977. A prokaryotic intracellular symbiont of the Great Salt Lake brine shrimp *Artemia salina* (L.). *Can. J. Microbiol.* 23: 1232-1236.
- Rodríguez-Almaraz, A.G., C. Zavala, R. Mendoza, & A.M. Maeda-Martínez. 2006. Ecological and biological notes on the brine shrimp *Artemia* (crustacean: Branchiopoda: Anostraca) from Carmen Island, Baja California Sur, México. *Hydrobiologia* 560: 417-423.
- Sokal, R.R. & J.F. Rohlf. 1981. *Biometry*. W.H. Freeman. San Francisco, California, USA.
- Tackaert, W. & P. Sorgeloos. 1991. Semi-intensive culturing in fertilized ponds, p. 287-312. *In* R.A. Browne, P. Sorgeloos & C.N.A. Trotman (eds.). *Artemia* Biology. CRC, Boston, USA.
- Tatsuoka, M.M. 1970. Selected topics in advanced statistics. An Elementary Approach. No. 6. Discriminant Analysis. IPAT, Illinois, USA.
- Triantaphyllidis, G.V., K. Pouloupoulou, T.J. Abatzopoulos, C.A. Pinto-Perez & P. Sorgeloos. 1995. International study on *Artemia* XLIX. Salinity effects on survival, maturity, growth, biometrics, reproductive and lifespan characteristics of a bisexual and a parthenogenetic population of *Artemia*. *Hydrobiologia* 302: 215-227.
- Triantaphyllidis, G.V., G.R.J. Criel, T.J. Abatzopoulos & P. Sorgeloos. 1997a. International study on *Artemia*. LIII. Morphological study of *Artemia* with emphasis to Old World populations. I. Bisexual populations. *Hydrobiologia* 357: 139-153.
- Triantaphyllidis, G.V., G.R.J. Criel, T.J. Abatzopoulos, K.M. Thomas, J. Peleman, J.A. Beardmore & P. Sorgeloos. 1997b. International study on *Artemia*. LVII. Morphological and molecular characters suggest conspecificity of all bisexual European and North African *Artemia* populations. *Mar. Biol.* 129: 477-487.
- Triantaphyllidis, G.V., T.J. Abatzopoulos & P. Sorgeloos. 1998. Review of the biogeography of the genus *Artemia* (Crustacea, Anostraca). *J. Biogeogr.* 25: 213-226.
- Vanhaecke, P., S.E. Siddal & P. Sorgeloos. 1984. International study on *Artemia*. XXII. Combined effects of temperature and salinity on the survival of *Artemia* of various geographical origins. *J. Exp. Mar. Biol. Ecol.* 80: 259-275.
- Van Stappen, G. 2002. Zoogeography, p. 171-224. *In* T.J. Abatzopoulos, J.A. Beardmore, J.S. Clegg & P. Sorgeloos (eds.), *Artemia: Basic and Applied Biology*. Kluwer Academic, Dordrecht, The Netherlands.
- Wear, R.G. & S.J. Haslett. 1986. Effects of temperature and salinity on the biology of *Artemia franciscana* Kellogg from Lake Grassmere, New Zealand. 1. Growth and mortality. *J. Exp. Mar. Biol. Ecol.* 98: 153-166.